

Azithromycin

FORM 200mg/5ml Oral Suspension (Dry powder for reconstitution)

INDICATION 2nd line treatment of ocular chlamydia trachomatis infection

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
Birth – 6 months	20mg/kg	Once daily for 3 days	Oral

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

- See Summary of Product Characteristics and most recent edition of BNF for Children (links below)

FURTHER INFORMATION

- Reconstitute as per manufacturer instructions on bottle

STORAGE

Store at room temperature after opening. Discard 10 days after reconstitution

LICENSED STATUS

Licensed for use under 15kg

LINKS

[BNF for Children:](#) / [Electronic Medicines Compendium](#)

APPLICABLE POLICIES

[West of Scotland Neonatal Guidelines:](#)

Consult local policy if applicable

Document Number:	001	Supersedes:	Nil
Prepared by:	Fiona Johnston	Checked by	WoS neonatal Pharmacists
Date prepared	February 2020	Date updated	
Updated by		Review Date	February 2023

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications. This monograph should be used in conjunction with the package insert, BNF for Children, and Summary of Product Characteristics. For further advice contact your clinical pharmacist or pharmacy department.