# **West of Scotland NEONATAL Parenteral Drug Monographs**

# **Hepatitis B Vaccine Engerix**

Please remember to complete all relevant local paperwork when giving Hepatitis B Vaccine / Immunoglobulin. This will ensure appropriate follow up in Primary Care (See Immunisation Protocol for more details)

BRAND NAME Engerix B<sup>®</sup>

(Note only this brand should be used with this monograph)

**FORM** Hepatitis B surface antigen 20micrograms/ml (0.5ml prefilled syringe)

INDICATION

Immunisation of babies born to mothers positive for Hepatitis B or in high risk groups for developing Hepatitis B (See Immunisation Policy for full

details).

### **DOSE RANGE**

AGE	DOSE	FREQUENCY	ROUTE
Neonate – 6 months	10micrograms (0.5ml)	Single dose	IM
		(See Vaccination schedule)	

#### **ADMINISTRATION**

Give by IM injection into anterolateral thigh.

Local reactions may be reduced by ensuring the needle is placed deep into the muscle.

If immunoglobulin is to be administered at the same time, inject at different sites.

## **CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS**

- See Summary of Product Characteristics and most recent edition of BNF for Children (links below)

#### **FURTHER INFORMATION**

Preterm babies follow this protocol with no adjustment for prematurity.

STORAGE Store in the fridge

LICENSED STATUS Licensed use

LINKS BNF for Children: / Electronic Medicines Compendium

APPLICABLE POLICIES West of Scotland Neonatal Guidelines:

Consult local policy if applicable

Document Number:	003	Supersedes:	002
Prepared by:	WoS Neo pharm group	Checked by	F. Anderson
Date prepared	May 2016	Date updated	April 2018
Updated by	June Grant	Review Date	April 2021

# Administer reconstituted solutions immediately.

All vials, ampoules and infusion bags are for single use only unless otherwise stated.

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications. This monograph should be used in conjunction with the package insert, BNF for Children, and Summary of Product Characteristics. For further advice contact your clinical pharmacist or pharmacy department.

Meomatall