

Pneumococcal Conjugated Vaccine, Adsorbed

Neonatal

BRAND Prevenar 13

FORM Pneumococcal conjugated vaccine, 0.5ml prefilled syringe

INDICATION Primary Immunisation (See WoS Immunisation Guideline for full details of immunisation programme)

DOSE RANGE

AGE	DOSE	FREQUENCY	ROUTE
2 - 6 months	0.5ml	ONCE only (repeat as per immunisation programme)	IM injection

ADMINISTRATION

Shake well to mix white deposit and clear supernatant

Give by IM injection into anterolateral thigh.

Local reactions may be reduced by ensuring the needle is placed deep into the muscle.

Injection should be given into a different limb to vaccines given at same time or at least 2.5cm apart if same limb MUST be used

CAUTIONS, CONTRA-INDICATIONS AND SIDE EFFECTS

- See Summary of Product Characteristics and most recent edition of BNF for Children (links below)

FURTHER INFORMATION

- Premature babies follow the same programme with no correction for gestational age
- Complete relevant local paperwork for public health
- The first course of vaccination has been associated with a recurrence of apnoea in premature babies. Such babies should be monitored closely following vaccination.

STORAGE Store in the fridge

LICENSED STATUS Licensed Use

LINKS [BNF for Children](#) / [Electronic Medicines Compendium](#)

APPLICABLE POLICIES [West of Scotland Neonatal Guidelines](#):

Consult local policy if applicable

Document Number:	002	Supersedes:	001
Prepared by:	WoS Neo pharm group	Checked by	F.Anderson
Date prepared	May 2016	Date updated	August 2019
Updated by	June Grant	Review Date	August 2022

Dose may vary depending on indication, age, renal function, hepatic function, and concomitant medications. This monograph should be used in conjunction with the package insert, BNF for Children, and Summary of Product Characteristics. For further advice contact your clinical pharmacist or pharmacy department.